arXiv:1003.3297v1 [math.NT] 17 Mar 2010

IDENTITIES OF SYMMETRY FOR q-BERNOULLI POLYNOMIALS

DAE SAN KIM

Abstract. In this paper, we derive eight basic identities of symmetry in three variables related to q-Bernoulli polynomials and the q-analogue of power sums. These and most of their corollaries are new, since there have been results only about identities of symmetry in two variables. These abundance of symmetries shed new light even on the existing identities so as to yield some further interesting ones. The derivations of identities are based on the p-adic integral expression of the generating function for the q-Bernoulli polynomials and the quotient of integrals that can be expressed as the exponential generating function for the q-analogue of power sums.

Key words : q-Bernoulli polynomial, q-analogue of power sum, Volkenborn integral, identities of symmetry.

MSC2010:11B68;11S80;05A19.

1. Introduction and preliminaries

Let p be a fixed prime. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}_p$ will respectively denote the ring of p-adic integers, the field of p-adic rational numbers and the completion of the algebraic closure of $\mathbb{Q}_p$. For a uniformly differentiable(also called continuously differentiable) function $f : \mathbb{Z}_p \longrightarrow \mathbb{C}_p$ (cf. [6]), the Volkenborn integral of f is defined by

$$\int_{\mathbb{Z}_p} f(z)d\mu(z) = \lim_{N\to\infty} \frac{1}{p^N} \sum_{j=0}^{p^N-1} f(j).$$

Then it is easy to see that

$$\int_{\mathbb{Z}_p} f(z+1)d\mu(z) = \int_{\mathbb{Z}_p} f(z)d\mu(z) + f'(0). \tag{1.1}$$

Let $|\ |_p$ be the normalized absolute value of $\mathbb{C}_p$, such that $|p|_p = \frac{1}{p}$, and let

$$E = \{t \in \mathbb{C}_p \,|\, |t|_p < p^{\frac{-1}{p-1}}\}. \tag{1.2}$$

Assume that $q, t \in \mathbb{C}_p$, with $q-1, t \in E$, so that $q^z = \exp(z \log q)$ and e^{zt} are, as functions of z, analytic functions on $\mathbb{Z}_p$. By applying (1.1) to f, with $f(z) = q^z e^{tz}$, we get the p-adic integral expression of the generating function for q-Bernoulli numbers $B_{n,q}$ (cf. [5], [7]):

$$\int_{\mathbb{Z}_p} q^z e^{zt} d\mu(z) = \frac{\log q + t}{qe^t - 1} = \sum_{n=0}^{\infty} B_{n,q} \frac{t^n}{n!} \ (q-1, t \in E). \tag{1.3}$$

So we have the following p-adic integral expression of the generating function for the q-Bernoulli polynomials $B_{n,q}(x)$:

$$\int_{\mathbb{Z}_p} q^z e^{(x+z)t} d\mu(z) = \frac{\log q + t}{qe^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_{n,q}(x) \frac{t^n}{n!} \ (q-1, t \in E, x \in \mathbb{Z}_p). \tag{1.4}$$

Here and throughout this paper, we will have many instances to be able to interchange integral and infinite sum. That is justified by Proposition 55.4 in [6]. Let $S_{k,q}(n)$ denote the q-analogue of kth power sum of the first n+1 nonnegative integers, namely

$$S_{k,q}(n) = \sum_{i=0}^{n} i^k q^i = 0^k q^0 + 1^k q^1 + \cdots + n^k q^n. \tag{1.5}$$

In particular,

$$S_{0,q}(n) = \sum_{i=0}^{n} q^i = \frac{q^{n+1}-1}{q-1} = [n+1]_q, \ S_{k,q}(0) = \begin{cases} 1, & \text{for } k = 0, \\ 0, & \text{for } k > 0. \end{cases} \tag{1.6}$$

From 1.3 and 1.5, one easily derives the following identities: for $w \in \mathbb{Z}_{>0}$,

$$\frac{w \int_{\mathbb{Z}_p} q^x e^{xt} d\mu(x)}{\int_{\mathbb{Z}_p} q^{wy} e^{wyt} d\mu(y)} = \sum_{i=0}^{w-1} q^i e^{it} = \sum_{k=0}^{\infty} S_{k,q}(w-1) \frac{t^k}{k!} \ (q-1, \ t \in E). \tag{1.7}$$

In what follows, we will always assume that the Volkenborn integrals of the various exponential functions on $\mathbb{Z}_p$ are defined for $q-1, t \in E$(cf. (1.2)), and therefore it will not be mentioned.

[1], [2], [4], [8] and [9] are some of the previous works on identities of symmetry in two variables involving Bernoulli polynomials and power sums. For the brief history, one is referred to those papers. In [3], the idea of [4] was adopted to produce many new identities of symmetry in three variables involving Bernoulli polynomials and power sums.

In this paper, we will produce 8 basic identities of symmetry in three variables w_1,w_2,w_3 related to q-Bernoulli polynomials and the q-analogue of power sums(cf. (4.8), (4.2), (4.12), (4.15), (4.19), (4.21), (4.23), (4.24)). These and most of their corollaries seem to be new, since there have been results only about identities of symmetry in two variables in the literature. These abundance of symmetries shed new light even on the existing identities. For instance, it has been known that (1.8) and (1.9) are equal(cf. [5, (2.20)]) and (1.10) and (1.11) are so(cf. [5, (2.25)]). In fact, (1.8)-(1.11) are all equal, as they can be derived from one and the same p-adic integral. Perhaps, this was neglected to mention in [5]. Also, we have a bunch of new identities in (1.12)-(1.15). All of these were obtained as corollaries(cf. Cor. 4.9, 4.12, 4.15) to some of the basic identities by specializing the variable w_3 as 1. Those would not be unearthed if more symmetries had not been available.

(1.8)
$$\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_2}}(w_1y_1)S_{n-k,q^{w_1}}(w_2-1)w_1^{n-k}w_2^{k-1}$$
(1.9)
$$=\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_1}}(w_2y_1)S_{n-k,q^{w_2}}(w_1-1)w_2^{n-k}w_1^{k-1}$$
(1.10)
$$=w_1^{n-1}\sum_{i=0}^{w_1-1}q^{w_2i}B_{n,q^{w_1}}(w_2y_1+\frac{w_2}{w_1}i)$$
(1.11)
$$=w_2^{n-1}\sum_{i=0}^{w_2-1}q^{w_1i}B_{n,q^{w_2}}(w_1y_1+\frac{w_1}{w_2}i)$$
(1.12)
$$=\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,q^{w_1w_2}}(y_1)S_{l,q^{w_2}}(w_1-1)S_{m,q^{w_1}}(w_2-1)w_1^{k+m-1}w_2^{k+l-1}$$
(1.13)
$$=w_1^{n-1}\sum_{k=0}^{n}\binom{n}{k}S_{n-k,q^{w_1}}(w_2-1)w_2^{k-1}\sum_{i=0}^{w_1-1}q^{w_2i}B_{k,q^{w_1w_2}}(y_1+\frac{i}{w_1})$$
(1.14)
$$=w_2^{n-1}\sum_{k=0}^{n}\binom{n}{k}S_{n-k,q^{w_2}}(w_1-1)w_1^{k-1}\sum_{i=0}^{w_2-1}q^{w_1i}B_{k,q^{w_1w_2}}(y_1+\frac{i}{w_2})$$
(1.15)
$$=(w_1w_2)^{n-1}\sum_{i=0}^{w_1-1}\sum_{j=0}^{w_2-1}q^{w_2i+w_1j}B_{n,q^{w_1w_2}}(y_1+\frac{i}{w_1}+\frac{j}{w_2}).$$

The derivations of identities are based on the p-adic integral expression of the generating function for the q-Bernoulli polynomials in (1.4) and the quotient of integrals in (1.7) that can be expressed as the exponential generating function for the q-analogue of power sums. We indebted this idea to the paper [5].

2. Several types of quotients of Volkenborn integrals

Here we will introduce several types of quotients of Volkenborn integrals on $\mathbb{Z}_p$ or $\mathbb{Z}_p^3$ from which some interesting identities follow owing to the built-in symmetries in w_1, w_2, w_3. In the following, w_1, w_2, w_3 are all positive integers and all of the explicit expressions of integrals in (2.2), (2.4), (2.6), and (2.8) are obtained from the identity in (1.3).

(a) Type Λ_{23}^{i} (for $i = 0, 1, 2, 3$)

$I(\Lambda_{23}^i)$

$$= \frac{\int_{\mathbb{Z}_p^3} q^{w_2w_3x_1+w_1w_3x_2+w_1w_2x_3} e^{(w_2w_3x_1+w_1w_3x_2+w_1w_2x_3+w_1w_2w_3(\sum_{j=1}^{3-i} y_j))t} d\mu(X)}{(\int_{\mathbb{Z}_p} q^{w_1w_2w_3x_4} e^{w_1w_2w_3x_4t} d\mu(x_4))^i}, \tag{2.1}$$

with $d\mu(X) = d\mu(x_1)d\mu(x_2)d\mu(x_3)$

$$= \frac{(w_1w_2w_3)^{2-i}(\log q + t)^{3-i} e^{w_1w_2w_3(\sum_{j=1}^{3-i} y_j)t}(q^{w_1w_2w_3}e^{w_1w_2w_3t}-1)^i}{(q^{w_2w_3}e^{w_2w_3t}-1)(q^{w_1w_3}e^{w_1w_3t}-1)(q^{w_1w_2}e^{w_1w_2t}-1)}; \tag{2.2}$$

(b) Type Λ_{13}^i (for $i = 0, 1, 2, 3$)
$I(\Lambda_{13}^i)$

$$= \frac{\int_{\mathbb{Z}_p^3} q^{w_1x_1+w_2x_2+w_3x_3} e^{(w_1x_1+w_2x_2+w_3x_3+w_1w_2w_3(\sum_{j=1}^{3-i} y_j))t} d\mu(x_1)d\mu(x_2)d\mu(x_3)}{(\int_{\mathbb{Z}_p} q^{w_1w_2w_3} e^{w_1w_2w_3x_4t} d\mu(x_4))^i} \tag{2.3}$$

$$= \frac{(w_1w_2w_3)^{1-i}(\log q + t)^{3-i} e^{w_1w_2w_3(\sum_{j=1}^{3-i} y_j)t}(q^{w_1w_2w_3}e^{w_1w_2w_3t}-1)^i}{(q^{w_1}e^{w_1t}-1)(q^{w_2}e^{w_2t}-1)(q^{w_3}e^{w_3t}-1)}; \tag{2.4}$$

(c-0) Type Λ_{12}^0
$I(\Lambda_{12}^0)$

$$= \int_{\mathbb{Z}_p^3} q^{w_1x_1+w_2x_2+w_3x_3} e^{(w_1x_1+w_2x_2+w_3x_3+w_2w_3y+w_1w_3y+w_1w_2y)t} d\mu(x_1)d\mu(x_2)d\mu(x_3) \tag{2.5}$$

$$= \frac{w_1w_2w_3(\log q + t)^3 e^{(w_2w_3+w_1w_3+w_1w_2)yt}}{(q^{w_1}e^{w_1t}-1)(q^{w_2}e^{w_2t}-1)(q^{w_3}e^{w_3t}-1)}; \tag{2.6}$$

(c-1) Type Λ_{12}^1

$$I(\Lambda_{12}^1) = \frac{\int_{\mathbb{Z}_p^3} q^{w_1x_1+w_2x_2+w_3x_3} e^{(w_1x_1+w_2x_2+w_3x_3)t} d\mu(x_1)d\mu(x_2)d\mu(x_3)}{\int_{\mathbb{Z}_p^3} q^{w_2w_3z_1+w_1w_3z_2+w_1w_2z_3} e^{(w_2w_3z_1+w_1w_3z_2+w_1w_2z_3)t} d\mu(z_1)d\mu(z_2)d\mu(z_3)} \tag{2.7}$$

$$= \frac{(w_1w_2w_3)^{-1}(q^{w_2w_3}e^{w_2w_3t}-1)(q^{w_1w_3}e^{w_1w_3t}-1)(q^{w_1w_2}e^{w_1w_2t}-1)}{(q^{w_1}e^{w_1t}-1)(q^{w_2}e^{w_2t}-1)(q^{w_3}e^{w_3t}-1)}. \tag{2.8}$$

All of the above p-adic integrals of various types are invariant under all permutations of w_1, w_2, w_3, as one can see either from p-adic integral representations in (2.1), (2.3), (2.5), and (2.7) or from their explicit evaluations in (2.2), (2.4), (2.6), and (2.8).

3. Identities for q-Bernoulli polynomials

First, let's consider Type Λ_{23}^{i}, for each $i = 0, 1, 2, 3$.
(a-0) The following results can be easily obtained from (1.4) and (1.7).

$$
\begin{aligned}
I(\Lambda_{23}^{0}) &= \int_{\mathbb{Z}_p} q^{w_2w_3x_1} e^{w_2w_3(x_1+w_1y_1)t} d\mu(x_1) \int_{\mathbb{Z}_p} q^{w_1w_3x_2} e^{w_1w_3(x_2+w_2y_2)t} d\mu(x_2) \\
&\qquad \times \int_{\mathbb{Z}_p} q^{w_1w_2x_3} e^{w_1w_2(x_3+w_3y_3)t} d\mu(x_3) \\
&= \Big(\sum_{k=0}^{\infty} \frac{B_{k,q^{w_2w_3}}(w_1y_1)}{k!} (w_2w_3t)^k\Big)\Big(\sum_{l=0}^{\infty} \frac{B_{l,q^{w_1w_3}}(w_2y_2)}{l!} (w_1w_3t)^l\Big) \\
&\qquad \times \Big(\sum_{m=0}^{\infty} \frac{B_{m,q^{w_1w_2}}(w_3y_3)}{m!} (w_1w_2t)^m\Big) \\
&= \sum_{n=0}^{\infty}\Big(\sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_2w_3}}(w_1y_1) B_{l,q^{w_1w_3}}(w_2y_2) \\
&\qquad \times B_{m,q^{w_1w_2}}(w_3y_3) w_1^{l+m} w_2^{k+m} w_3^{k+l}\Big) \frac{t^n}{n!},
\end{aligned}
\tag{3.1}
$$

where the inner sum is over all nonnegative integers k, l, m, with $k + l + m = n$, and

$$
\binom{n}{k,l,m} = \frac{n!}{k!l!m!}.
\tag{3.2}
$$

(a-1) Here we write $I(\Lambda_{23}^{1})$ in two different ways:

(1) $I(\Lambda_{23}^{1})$

$$
\begin{aligned}
&= \frac{1}{w_3} \int_{\mathbb{Z}_p} q^{w_2w_3x_1} e^{w_2w_3(x_1+w_1y_1)t} d\mu(x_1) \int_{\mathbb{Z}_p} q^{w_1w_3x_2} e^{w_1w_3(x_2+w_2y_2)t} d\mu(x_2) \\
&\qquad \times \frac{w_3 \int_{\mathbb{Z}_p} q^{w_1w_2x_3} e^{w_1w_2x_3t} d\mu(x_3)}{\int_{\mathbb{Z}_p} q^{w_1w_2w_3x_4} e^{w_1w_2w_3x_4t} d\mu(x_4)}
\end{aligned}
\tag{3.3}
$$

$$
\begin{aligned}
&= \frac{1}{w_3}\Big(\sum_{k=0}^{\infty} B_{k,q^{w_2w_3}}(w_1y_1) \frac{(w_2w_3t)^k}{k!}\Big) \sum_{l=0}^{\infty} B_{l,q^{w_1w_3}}(w_2y_2) \frac{(w_1w_3t)^l}{l!}\Big) \\
&\qquad \times \Big(\sum_{m=0}^{\infty} S_{m,q^{w_1w_2}}(w_3-1) \frac{(w_1w_2t)^m}{m!}\Big)
\end{aligned}
$$

$$
\begin{aligned}
&= \sum_{n=0}^{\infty}\Big(\sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_2w_3}}(w_1y_1) B_{l,q^{w_1w_3}}(w_2y_2) \\
&\qquad \times S_{m,q^{w_1w_2}}(w_3-1) w_1^{l+m} w_2^{k+m} w_3^{k+l-1}\Big) \frac{t^n}{n!}.
\end{aligned}
\tag{3.4}
$$

(2) Invoking (1.7), (3.3) can also be written as

$$\begin{aligned}
&I(\Lambda_{23}^{1})\\
&=\frac{1}{w_3}\sum_{i=0}^{w_3-1}q^{w_1w_2i}\int_{\mathbb{Z}_p}q^{w_2w_3x_1}e^{w_2w_3(x_1+w_1y_1)t}d\mu(x_1)\\
&\qquad\qquad\times\int_{\mathbb{Z}_p}q^{w_1w_3x_2}e^{w_1w_3(x_2+w_2y_2+\frac{w_2}{w_3}i)t}d\mu(x_2)\\
&=\frac{1}{w_3}\sum_{i=0}^{w_3-1}q^{w_1w_2i}(\sum_{k=0}^{\infty}B_{k,q^{w_2w_3}}(w_1y_1)\frac{(w_2w_3t)^k}{k!})(\sum_{l=0}^{\infty}B_{l,q^{w_1w_3}}(w_2y_2+\frac{w_2}{w_3}i)\frac{(w_1w_3t)^l}{l!})\\
&=\sum_{n=0}^{\infty}(w_3^{n-1}\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_2w_3}}(w_1y_1)w_1^{n-k}w_2^{k}\sum_{i=0}^{w_3-1}q^{w_1w_2i}B_{n-k,q^{w_1w_3}}(w_2y_2+\frac{w_2}{w_3}i))\frac{t^n}{n!}.
\end{aligned}\tag{3.5}$$

(a-2) Here we write $I(\Lambda_{23}^{2})$ in three different ways:

(1) $I(\Lambda_{23}^{2})$

$$\begin{aligned}
&=\frac{1}{w_2w_3}\int_{\mathbb{Z}_p}q^{w_2w_3x_1}e^{w_2w_3(x_1+w_1y_1)t}d\mu(x_1)\\
&\qquad\times\frac{w_2\int_{\mathbb{Z}_p}q^{w_1w_3x_2}e^{w_1w_3x_2t}d\mu(x_2)}{\int_{\mathbb{Z}_p}q^{w_1w_2w_3x_4}e^{w_1w_2w_3x_4t}d\mu(x_4)}\times\frac{w_3\int_{\mathbb{Z}_p}q^{w_1w_2x_3}e^{w_1w_2x_3t}d\mu(x_3)}{\int_{\mathbb{Z}_p}q^{w_1w_2w_3x_4}e^{w_1w_2w_3x_4t}d\mu(x_4)}
\end{aligned}\tag{3.6}$$

$$\begin{aligned}
&=\frac{1}{w_2w_3}(\sum_{k=0}^{\infty}B_{k,q^{w_2w_3}}(w_1y_1)\frac{(w_2w_3t)^k}{k!})\\
&\qquad\times(\sum_{l=0}^{\infty}S_{l,q^{w_1w_3}}(w_2-1)\frac{(w_1w_3t)^l}{l!})(\sum_{m=0}^{\infty}S_{m,q^{w_1w_2}}(w_3-1)\frac{(w_1w_2t)^m}{m!})\\
&=\sum_{n=0}^{\infty}(\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,q^{w_2w_3}}(w_1y_1)S_{l,q^{w_1w_3}}(w_2-1)S_{m,q^{w_1w_2}}(w_3-1)\\
&\qquad\qquad\times w_1^{l+m}w_2^{k+m-1}w_3^{k+l-1})\frac{t^n}{n!}.
\end{aligned}\tag{3.7}$$

(2) Invoking (1.7), (3.6) can also be written as

$I(\Lambda_{23}^{2})$

$$\begin{aligned}
&=\frac{1}{w_2w_3}\sum_{i=0}^{w_2-1}q^{w_1w_3i}\int_{\mathbb{Z}_p}q^{w_2w_3x_1}e^{w_2w_3(x_1+w_1y_1+\frac{w_1}{w_2}i)t}d\mu(x_1)\\
&\qquad\times\frac{w_2\int_{\mathbb{Z}_p}q^{w_1w_2x_3}e^{w_1w_2x_3t}d\mu(x_3)}{\int_{\mathbb{Z}_p}q^{w_1w_2w_3x_4}e^{w_1w_2w_3x_4t}d\mu(x_4)}
\end{aligned}\tag{3.8}$$

$$\begin{aligned}
&= \frac{1}{w_2w_3}\sum_{i=0}^{w_2-1} q^{w_1w_3i}\Big(\sum_{k=0}^{\infty} B_{k,q^{w_2w_3}}\big(w_1y_1+\frac{w_1}{w_2}i\big)\frac{(w_2w_3t)^k}{k!}\Big)\\
&\qquad\qquad\qquad\qquad \times\Big(\sum_{l=0}^{\infty} S_{l,q^{w_1w_2}}(w_3-1)\frac{(w_1w_2t)^l}{l!}\Big)\\
&= \sum_{n=0}^{\infty}\Big(w_2^{n-1}\sum_{k=0}^{n}\binom{n}{k}S_{n-k,q^{w_1w_2}}(w_3-1)w_1^{n-k}w_3^{k-1}\\
&\qquad\qquad\qquad\qquad \times\sum_{i=0}^{w_2-1} q^{w_1w_3i}B_{k,q^{w_2w_3}}\big(w_1y_1+\frac{w_1}{w_2}i\big)\Big)\frac{t^n}{n!}.
\end{aligned}\tag{3.9}$$

(3) Invoking (1.7) once again, (3.8) can be written as

$$\begin{aligned}
&I(\Lambda_{23}^{2})\\
&\quad= \frac{1}{w_2w_3}\sum_{i=0}^{w_2-1}\sum_{j=0}^{w_3-1} q^{w_1w_3i+w_1w_2j}\int_{\mathbb{Z}_p} q^{w_2w_3x_1}e^{w_2w_3(x_1+w_1y_1+\frac{w_1}{w_2}i+\frac{w_1}{w_3}j)t}d\mu(x_1)\\
&\quad= \frac{1}{w_2w_3}\sum_{i=0}^{w_2-1}\sum_{j=0}^{w_3-1}\Big(q^{w_1w_3i+w_1w_2j}\sum_{n=0}^{\infty}B_{n,q^{w_2w_3}}\big(w_1y_1+\frac{w_1}{w_2}i+\frac{w_1}{w_3}j\big)\frac{(w_2w_3t)^n}{n!}\Big)\\
&\quad= \sum_{n=0}^{\infty}\Big((w_2w_3)^{n-1}\sum_{i=0}^{w_2-1}\sum_{j=0}^{w_3-1}q^{w_1w_3i+w_1w_2j}B_{n,q^{w_2w_3}}\big(w_1y_1+\frac{w_1}{w_2}i+\frac{w_1}{w_3}j\big)\Big)\frac{t^n}{n!}.
\end{aligned}\tag{3.10}$$

(a-3)

$$\begin{aligned}
&I(\Lambda_{23}^{3})\\
&\quad= \frac{1}{w_1w_2w_3}\frac{w_1\int_{\mathbb{Z}_p} q^{w_2w_3x_1}e^{w_2w_3x_1t}d\mu(x_1)}{\int_{\mathbb{Z}_p} q^{w_1w_2w_3x_4}e^{w_1w_2w_3x_4t}d\mu(x_4)}\\
&\qquad\qquad \times\frac{w_2\int_{\mathbb{Z}_p} q^{w_1w_3x_2}e^{w_1w_3x_2t}d\mu(x_2)}{\int_{\mathbb{Z}_p} q^{w_1w_2w_3x_4}e^{w_1w_2w_3x_4t}d\mu(x_4)}\times\frac{w_3\int_{\mathbb{Z}_p} q^{w_1w_2x_3}e^{w_1w_2x_3t}d\mu(x_3)}{\int_{\mathbb{Z}_p} q^{w_1w_2w_3x_4}e^{w_1w_2w_3x_4t}d\mu(x_4)}\\
&\quad= \frac{1}{w_1w_2w_3}\Big(\sum_{k=0}^{\infty}S_{k,q^{w_2w_3}}(w_1-1)\frac{(w_2w_3t)^k}{k!}\Big)\\
&\qquad\qquad \times\Big(\sum_{l=0}^{\infty}S_{l,q^{w_1w_3}}(w_2-1)\frac{(w_1w_3t)^l}{l!}\Big)\Big(\sum_{m=0}^{\infty}S_{m,q^{w_1w_2}}(w_3-1)\frac{(w_1w_2t)^m}{m!}\Big)\\
&\quad= \sum_{n=0}^{\infty}\Big(\sum_{k+l+m=n}\binom{n}{k,l,m}S_{k,q^{w_2w_3}}(w_1-1)S_{l,q^{w_1w_3}}(w_2-1)S_{m,q^{w_1w_2}}(w_3-1)\\
&\qquad\qquad\qquad\qquad \times w_1^{l+m-1}w_2^{k+m-1}w_3^{k+l-1}\Big)\frac{t^n}{n!}.
\end{aligned}\tag{3.11}$$

(b) For Type Λ_{13}^{i} $(i=0,1,2,3)$, we may consider the analogous things to the ones in (a-0), (a-1), (a-2), and (a-3). However, these do not lead us to new identities. Indeed, if we substitute w_2w_3, w_1w_3, w_1w_2 respectively for w_1, w_2, w_3 in (2.1), this amounts to replacing t by $w_1w_2w_3t$ and q by $q^{w_1w_2w_3}$ in (2.3). So, upon replacing w_1, w_2, w_3 respectively by w_2w_3, w_1w_3, w_1w_2 and dividing by $(w_1w_2w_3)^n$, in each of the expressions of (3.1),(3.4), (3.5), (3.7), (3.9)-(3.11), we will get the corresponding symmetric identities for Type Λ_{13}^{i} $(i=0,1,2,3)$.

(c-0)

$$\begin{aligned}
&I(\Lambda_{12}^{0})\\
&\quad=\int_{\mathbb{Z}_p} q^{w_1x_1}e^{w_1(x_1+w_2y)t}d\mu(x_1)\\
&\qquad\qquad\times\int_{\mathbb{Z}_p} q^{w_2x_2}e^{w_2(x_2+w_3y)t}d\mu(x_2)\int_{\mathbb{Z}_p} q^{w_3x_3}e^{w_3(x_3+w_1y)t}d\mu(x_3)\\
&\quad=(\sum_{n=0}^{\infty}\frac{B_{k,q^{w_1}}(w_2y)}{k!}(w_1t)^k)(\sum_{l=0}^{\infty}\frac{B_{l,q^{w_2}}(w_3y)}{l!}(w_2t)^l)(\sum_{m=0}^{\infty}\frac{B_{m,q^{w_3}}(w_1y)}{m!}(w_3t)^m)
\end{aligned}$$

(3.12)

$$=\sum_{n=0}^{\infty}(\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,q^{w_1}}(w_2y)B_{l,q^{w_2}}(w_3y)B_{m,q^{w_3}}(w_1y)w_1^kw_2^lw_3^m)\frac{t^n}{n!}.$$

(c-1)

$$\begin{aligned}
I(\Lambda_{12}^{1})&=\frac{1}{w_1w_2w_3}\frac{w_2\int_{\mathbb{Z}_p}q^{w_1x_1}e^{w_1x_1t}d\mu(x_1)}{\int_{\mathbb{Z}_p}q^{w_1w_2z_3}e^{w_1w_2z_3t}d\mu(z_3)}\\
&\qquad\times\frac{w_3\int_{\mathbb{Z}_p}q^{w_2x_2}e^{w_2x_2t}d\mu(x_2)}{\int_{\mathbb{Z}_p}q^{w_2w_3z_1}e^{w_2w_3z_1t}d\mu(z_1)}\times\frac{w_1\int_{\mathbb{Z}_p}q^{w_3x_3}e^{w_3x_3t}d\mu(x_3)}{\int_{\mathbb{Z}_p}q^{w_3w_1z_2}e^{w_3w_1z_2t}d\mu(z_2)}\\
&=\frac{1}{w_1w_2w_3}(\sum_{k=0}^{\infty}S_{k,q^{w_1}}(w_2-1)\frac{(w_1t)^k}{k!})\\
&\qquad\times(\sum_{l=0}^{\infty}S_{l,q^{w_2}}(w_3-1)\frac{(w_2t)^l}{l!})(\sum_{m=0}^{\infty}S_{m,q^{w_3}}(w_1-1)\frac{(w_3t)^m}{m!})\\
&=\sum_{n=0}^{\infty}(\sum_{k+l+m=n}\binom{n}{k,l,m}S_{k,q^{w_1}}(w_2-1)S_{l,q^{w_2}}(w_3-1)\\
&\qquad\times S_{m,q^{w_3}}(w_1-1)w_1^{k-1}w_2^{l-1}w_3^{m-1})\frac{t^n}{n!}.
\end{aligned}\tag{3.13}$$

4. Main theorems

As we noted earlier in the last paragraph of Section 2, the various types of quotients of Volkenborn integrals are invariant under any permutation of w_1, w_2, w_3. So the corresponding expressions in Section 3. are also invariant under any permutation of w_1, w_2, w_3. Thus our results about identities of symmetry will be immediate consequences of this observation.

However, not all permutations of an expression in Section 3 yield distinct ones. In fact, as these expressions are obtained by permuting w_1, w_2, w_3 in a single one labeled by them, they can be viewed as a group in a natural manner and hence it is isomorphic to a quotient of S_3. In particular, the number of possible distinct expressions are $1, 2, 3$, or 6. (a-0), (a-1(1)), (a-1(2)), and (a-2(2)) give the full six identities of symmetry, (a-2(1)) and (a-2(3)) yield three identities of symmetry, and (c-0) and (c-1) give two identities of symmetry, while the expression in (a-3) yields no identities of symmetry.

Here we will just consider the cases of Theorems 4.8 and 4.17, leaving the others as easy exercises for the reader. As for the case of Theorem 4.8, in addition to (4.14)-(4.16), we get the following three ones:

$$\sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_2 w_3}}(w_1 y_1) S_{l,q^{w_1 w_2}}(w_3-1) S_{m,q^{w_1 w_3}}(w_2-1) \times w_1^{l+m} w_3^{k+m-1} w_2^{k+l-1}, \tag{4.1}$$

$$\sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1 w_3}}(w_2 y_1) S_{l,q^{w_2 w_3}}(w_1-1) S_{m,q^{w_1 w_2}}(w_3-1) \times w_2^{l+m} w_1^{k+m-1} w_3^{k+l-1}, \tag{4.2}$$

$$\sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1 w_2}}(w_3 y_1) S_{l,q^{w_1 w_3}}(w_2-1) S_{m,q^{w_2 w_3}}(w_1-1) \times w_3^{l+m} w_2^{k+m-1} w_1^{k+l-1}. \tag{4.3}$$

But, by interchanging l and m, we see that (4.1), (4.2), and (4.3) are respectively equal to (4.14), (4.15), and (4.16).

As to Theorem 4.17, in addition to (4.24) and (4.25), we have:

$$\sum_{k+l+m=n} \binom{n}{k,l,m} S_{k,q^{w_1}}(w_2-1) S_{l,q^{w_2}}(w_3-1) S_{m,q^{w_3}}(w_1-1) w_1^{k-1} w_2^{l-1} w_3^{m-1}, \tag{4.4}$$

$$\sum_{k+l+m=n} \binom{n}{k,l,m} S_{k,q^{w_2}}(w_3-1) S_{l,q^{w_3}}(w_1-1) S_{m,q^{w_1}}(w_2-1) w_2^{k-1} w_3^{l-1} w_1^{m-1}, \tag{4.5}$$

$$\sum_{k+l+m=n} \binom{n}{k,l,m} S_{k,q^{w_1}}(w_3-1) S_{l,q^{w_3}}(w_2-1) S_{m,q^{w_2}}(w_1-1) w_1^{k-1} w_3^{l-1} w_2^{m-1}, \tag{4.6}$$

$$\sum_{k+l+m=n} \binom{n}{k,l,m} S_{k,q^{w_3}}(w_2-1) S_{l,q^{w_2}}(w_1-1) S_{m,q^{w_1}}(w_3-1) w_3^{k-1} w_2^{l-1} w_1^{m-1}. \tag{4.7}$$

However, (4.4) and (4.5) are equal to (4.24), as we can see by applying the permutations $k \to l, l \to m, m \to k$ for (4.4) and $k \to m, l \to k, m \to l$ for (4.5).

Similarly, we see that (4.6) and (4.7) are equal to (4.25), by applying permutations $k \to l, l \to m, m \to k$ for (4.6) and $k \to m, l \to k, m \to l$ for (4.7).

Theorem 4.1. *Let w_1, w_2, w_3 be any positive integers. Then the following expression is invariant under any permutation of w_1, w_2, w_3, so that it gives us six symmetries.*

$$\begin{aligned}
&\sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_2w_3}}(w_1y_1)B_{l,q^{w_1w_3}}(w_2y_2)B_{m,q^{w_1w_2}}(w_3y_3)w_1^{l+m}w_2^{k+m}w_3^{k+l} \\
&= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_2w_3}}(w_1y_1)B_{l,q^{w_1w_2}}(w_3y_2)B_{m,q^{w_1w_3}}(w_2y_3)w_1^{l+m}w_3^{k+m}w_2^{k+l} \\
&= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1w_3}}(w_2y_1)B_{l,q^{w_2w_3}}(w_1y_2)B_{m,q^{w_1w_2}}(w_3y_3)w_2^{l+m}w_1^{k+m}w_3^{k+l} \\
&= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1w_3}}(w_2y_1)B_{l,q^{w_1w_2}}(w_3y_2)B_{m,q^{w_2w_3}}(w_1y_3)w_2^{l+m}w_3^{k+m}w_1^{k+l} \\
&= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1w_2}}(w_3y_1)B_{l,q^{w_2w_3}}(w_1y_2)B_{m,q^{w_1w_3}}(w_2y_3)w_3^{l+m}w_1^{k+m}w_2^{k+l} \\
&= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1w_2}}(w_3y_1)B_{l,q^{w_1w_3}}(w_2y_2)B_{m,q^{w_2w_3}}(w_1y_3)w_3^{l+m}w_2^{k+m}w_1^{k+l}.
\end{aligned} \tag{4.8}$$

Theorem 4.2. *Let w_1, w_2, w_3 be any positive integers. Then the following expression is invariant under any permutation of w_1, w_2, w_3, so that it gives us six symmetries.*

$$\begin{aligned}
&\sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_2w_3}}(w_1y_1)B_{l,q^{w_1w_3}}(w_2y_2)S_{m,q^{w_1w_2}}(w_3-1) \\
&\qquad \times w_1^{l+m}w_2^{k+m}w_3^{k+l-1} \\
&= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_2w_3}}(w_1y_1)B_{l,q^{w_1w_2}}(w_3y_2)S_{m,q^{w_1w_3}}(w_2-1) \\
&\qquad \times w_1^{l+m}w_3^{k+m}w_2^{k+l-1} \\
&= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1w_3}}(w_2y_1)B_{l,q^{w_2w_3}}(w_1y_2)S_{m,q^{w_1w_2}}(w_3-1) \\
&\qquad \times w_2^{l+m}w_1^{k+m}w_3^{k+l-1} \\
&= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1w_3}}(w_2y_1)B_{l,q^{w_1w_2}}(w_3y_2)S_{m,q^{w_2w_3}}(w_1-1) \\
&\qquad \times w_2^{l+m}w_3^{k+m}w_1^{k+l-1} \\
&= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1w_2}}(w_3y_1)B_{l,q^{w_1w_3}}(w_2y_2)S_{m,q^{w_2w_3}}(w_1-1) \\
&\qquad \times w_3^{l+m}w_2^{k+m}w_1^{k+l-1}
\end{aligned} \tag{4.9}$$

$$\begin{aligned}
&= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1 w_2}}(w_3 y_1) B_{l,q^{w_2 w_3}}(w_1 y_2) S_{m,q^{w_1 w_3}}(w_2-1) \\
&\qquad\qquad\qquad\qquad \times w_3^{l+m} w_1^{k+m} w_2^{k+l-1}.
\end{aligned}$$

Putting $w_3 = 1$ in (4.2), we get the following corollary.

Corollary 4.3. *Let w_1, w_2 be any positive integers.*

$$\begin{aligned}
&\sum_{k=0}^{n} \binom{n}{k} B_{k,q^{w_2}}(w_1 y_1) B_{n-k,q^{w_1}}(w_2 y_2) w_1^{n-k} w_2^{k} \\
=&\sum_{k=0}^{n} \binom{n}{k} B_{k,q^{w_1}}(w_2 y_1) B_{n-k,q^{w_2}}(w_1 y_2) w_2^{n-k} w_1^{k} \\
=& \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1 w_2}}(y_1) B_{l,q^{w_1}}(w_2 y_2) S_{m,q^{w_2}}(w_1-1) w_2^{k+m} w_1^{k+l-1} \\
=& \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1}}(w_2 y_1) B_{l,q^{w_1 w_2}}(y_2) S_{m,q^{w_2}}(w_1-1) w_2^{l+m} w_1^{k+l-1} \\
=& \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1 w_2}}(y_1) B_{l,q^{w_2}}(w_1 y_2) S_{m,q^{w_1}}(w_2-1) w_1^{k+m} w_2^{k+l-1} \\
=& \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_2}}(w_1 y_1) B_{l,q^{w_1 w_2}}(y_2) S_{m,q^{w_1}}(w_2-1) w_1^{l+m} w_2^{k+l-1}.
\end{aligned} \tag{4.10}$$

Letting further $w_2 = 1$ in (4.10), we have the following corollary.

Corollary 4.4. *Let w_1 be any positive integer.*

$$\begin{aligned}
&\sum_{k=0}^{n} \binom{n}{k} B_{k,q}(w_1 y_1) B_{n-k,q^{w_1}}(y_2) w_1^{n-k} \\
=&\sum_{k=0}^{n} \binom{n}{k} B_{k,q^{w_1}}(y_1) B_{n-k,q}(w_1 y_2) w_1^{k} \\
=& \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1}}(y_1) B_{l,q^{w_1}}(y_2) S_{m,q}(w_1-1) w_1^{k+l-1}.
\end{aligned} \tag{4.11}$$

Theorem 4.5. *Let w_1, w_2, w_3 be any positive integers. Then the following expression is invariant under any permutation of w_1, w_2, w_3, so that it gives us six symmetries.*

(4.12)

$$\begin{aligned}
&w_1^{n-1} \sum_{k=0}^{n} \binom{n}{k} B_{k,q^{w_1 w_2}}(w_3 y_1) w_3^{n-k} w_2^{k} \sum_{i=0}^{w_1-1} q^{w_2 w_3 i} B_{n-k,q^{w_1 w_3}}(w_2 y_2 + \frac{w_2}{w_1} i) \\
=&w_1^{n-1} \sum_{k=0}^{n} \binom{n}{k} B_{k,q^{w_1 w_3}}(w_2 y_1) w_2^{n-k} w_3^{k} \sum_{i=0}^{w_1-1} q^{w_2 w_3 i} B_{n-k,q^{w_1 w_2}}(w_3 y_2 + \frac{w_3}{w_1} i)
\end{aligned}$$

$$\begin{aligned}
&=w_2^{n-1}\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_1w_2}}(w_3y_1)w_3^{n-k}w_1^{k}\sum_{i=0}^{w_2-1}q^{w_1w_3i}B_{n-k,q^{w_2w_3}}(w_1y_2+\frac{w_1}{w_2}i)\\
&=w_2^{n-1}\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_2w_3}}(w_1y_1)w_1^{n-k}w_3^{k}\sum_{i=0}^{w_2-1}q^{w_1w_3i}B_{n-k,q^{w_1w_2}}(w_3y_2+\frac{w_3}{w_2}i)\\
&=w_3^{n-1}\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_1w_3}}(w_2y_1)w_2^{n-k}w_1^{k}\sum_{i=0}^{w_3-1}q^{w_1w_2i}B_{n-k,q^{w_2w_3}}(w_1y_2+\frac{w_1}{w_3}i)\\
&=w_3^{n-1}\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_2w_3}}(w_1y_1)w_1^{n-k}w_2^{k}\sum_{i=0}^{w_3-1}q^{w_1w_2i}B_{n-k,q^{w_1w_3}}(w_2y_2+\frac{w_2}{w_3}i).
\end{aligned}$$

Letting $w_3=1$ in (4.12), we obtain alternative expressions for the identities in (4.10).

Corollary 4.6. *Let w_1, w_2 be any positive integers.*

$$\begin{aligned}
&\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_2}}(w_1y_1)B_{n-k,q^{w_1}}(w_2y_2)w_1^{n-k}w_2^{k}\\
&=\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_1}}(w_2y_1)B_{n-k,q^{w_2}}(w_1y_2)w_2^{n-k}w_1^{k}\\
&=w_1^{n-1}\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_1w_2}}(y_1)w_2^{k}\sum_{i=0}^{w_1-1}q^{w_2i}B_{n-k,q^{w_1}}(w_2y_2+\frac{w_2}{w_1}i)\\
&=w_1^{n-1}\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_1}}(w_2y_1)w_2^{n-k}\sum_{i=0}^{w_1-1}q^{w_2i}B_{n-k,q^{w_1w_2}}(y_2+\frac{i}{w_1})\\
&=w_2^{n-1}\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_1w_2}}(y_1)w_1^{k}\sum_{i=0}^{w_2-1}q^{w_1i}B_{n-k,q^{w_2}}(w_1y_2+\frac{w_1}{w_2}i)\\
&=w_2^{n-1}\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_2}}(w_1y_1)w_1^{n-k}\sum_{i=0}^{w_2-1}q^{w_1i}B_{n-k,q^{w_1w_2}}(y_2+\frac{i}{w_2}).
\end{aligned}\tag{4.13}$$

Putting further $w_2=1$ in (4.13), we have the alternative expressions for the identities for (4.11).

Corollary 4.7. *Let w_1 be any positive integer.*

$$\begin{aligned}
&\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_1}}(y_1)B_{n-k,q}(w_1y_2)w_1^{k}\\
&=\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_1}}(y_2)B_{n-k,q}(w_1y_1)w_1^{k}\\
&=w_1^{n-1}\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_1}}(y_1)\sum_{i=0}^{w_1-1}q^{i}B_{n-k,q^{w_1}}(y_2+\frac{i}{w_1}).
\end{aligned}$$

Theorem 4.8. *Let w_1, w_2, w_3 be any positive integers. Then we have the following three symmetries in w_1, w_2, w_3:*

$$\sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_2 w_3}}(w_1 y_1) S_{l,q^{w_1 w_3}}(w_2 - 1) S_{m,q^{w_1 w_2}}(w_3 - 1) \times w_1^{l+m} w_2^{k+m-1} w_3^{k+l-1} \tag{4.14}$$

$$= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1 w_3}}(w_2 y_1) S_{l,q^{w_1 w_2}}(w_3 - 1) S_{m,q^{w_2 w_3}}(w_1 - 1) \times w_2^{l+m} w_3^{k+m-1} w_1^{k+l-1} \tag{4.15}$$

$$= \sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1 w_2}}(w_3 y_1) S_{l,q^{w_2 w_3}}(w_1 - 1) S_{m,q^{w_1 w_3}}(w_2 - 1) \times w_3^{l+m} w_1^{k+m-1} w_2^{k+l-1}. \tag{4.16}$$

Putting $w_3 = 1$ in (4.14)-(4.16), we get the following corollary.

Corollary 4.9. *Let w_1, w_2 be any positive integers.*

$$\begin{aligned} &\sum_{k=0}^{n} \binom{n}{k} B_{k,q^{w_2}}(w_1 y_1) S_{n-k,q^{w_1}}(w_2 - 1) w_1^{n-k} w_2^{k-1} \\ =&\sum_{k=0}^{n} \binom{n}{k} B_{k,q^{w_1}}(w_2 y_1) S_{n-k,q^{w_2}}(w_1 - 1) w_2^{n-k} w_1^{k-1} \\ =&\sum_{k+l+m=n} \binom{n}{k,l,m} B_{k,q^{w_1 w_2}}(y_1) S_{l,q^{w_2}}(w_1 - 1) S_{m,q^{w_1}}(w_2 - 1) w_1^{k+m-1} w_2^{k+l-1}. \end{aligned} \tag{4.17}$$

Letting further $w_2 = 1$ in (4.17), we get the following corollary. This is also obtained in [5, (2.22)].

Corollary 4.10. *Let w_1 be any positive integer.*

$$B_{n,q}(w_1 y_1) = \sum_{k=0}^{n} \binom{n}{k} B_{k,q^{w_1}}(y_1) S_{n-k,q}(w_1 - 1) w_1^{k-1}. \tag{4.18}$$

Theorem 4.11. *Let w_1, w_2, w_3 be any positive integers. Then the following expression is invariant under any permutation of w_1, w_2, w_3, so that it gives us six symmetries.*

$$\begin{aligned} &w_1^{n-1} \sum_{k=0}^{n} \binom{n}{k} S_{n-k,q^{w_1 w_2}}(w_3 - 1) w_2^{n-k} w_3^{k-1} \sum_{i=0}^{w_1 - 1} q^{w_2 w_3 i} B_{k,q^{w_1 w_3}}(w_2 y_1 + \frac{w_2}{w_1} i) \\ =&w_1^{n-1} \sum_{k=0}^{n} \binom{n}{k} S_{n-k,q^{w_1 w_3}}(w_2 - 1) w_3^{n-k} w_2^{k-1} \sum_{i=0}^{w_1 - 1} q^{w_2 w_3 i} B_{k,q^{w_1 w_2}}(w_3 y_1 + \frac{w_3}{w_1} i) \end{aligned} \tag{4.19}$$

$$
\begin{aligned}
&=w_2^{n-1}\sum_{k=0}^{n}\binom{n}{k}S_{n-k,q^{w_1w_2}}(w_3-1)w_1^{n-k}w_3^{k-1}\sum_{i=0}^{w_2-1}q^{w_1w_3i}B_{k,q^{w_2w_3}}(w_1y_1+\frac{w_1}{w_2}i)\\
&=w_2^{n-1}\sum_{k=0}^{n}\binom{n}{k}S_{n-k,q^{w_2w_3}}(w_1-1)w_3^{n-k}w_1^{k-1}\sum_{i=0}^{w_2-1}q^{w_1w_3i}B_{k,q^{w_1w_2}}(w_3y_1+\frac{w_3}{w_1}i)\\
&=w_3^{n-1}\sum_{k=0}^{n}\binom{n}{k}S_{n-k,q^{w_1w_3}}(w_2-1)w_1^{n-k}w_2^{k-1}\sum_{i=0}^{w_3-1}q^{w_1w_2i}B_{k,q^{w_2w_3}}(w_1y_1+\frac{w_1}{w_3}i)\\
&=w_3^{n-1}\sum_{k=0}^{n}\binom{n}{k}S_{n-k,q^{w_2w_3}}(w_1-1)w_2^{n-k}w_1^{k-1}\sum_{i=0}^{w_3-1}q^{w_1w_2i}B_{k,q^{w_1w_3}}(w_2y_1+\frac{w_2}{w_3}i).
\end{aligned}
$$

Putting $w_3 = 1$ in (4.19), we obtain the following corollary. In Section 1, the identities in (4.17), (4.20), and (4.22) are combined to give those in (1.8)-(1.15).

Corollary 4.12. *Let w_1, w_2 be any positive integers.*

$$
\begin{aligned}
&w_1^{n-1}\sum_{i=0}^{w_1-1}q^{w_2i}B_{n,q^{w_1}}(w_2y_1+\frac{w_2}{w_1}i)\\
&=w_2^{n-1}\sum_{i=0}^{w_2-1}q^{w_1i}B_{n,q^{w_2}}(w_1y_1+\frac{w_1}{w_2}i)\\
&=\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_1}}(w_2y_1)S_{n-k,q^{w_2}}(w_1-1)w_2^{n-k}w_1^{k-1}\\
&=\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_2}}(w_1y_1)S_{n-k,q^{w_1}}(w_2-1)w_1^{n-k}w_2^{k-1}\\
&=w_1^{n-1}\sum_{k=0}^{n}\binom{n}{k}S_{n-k,q^{w_1}}(w_2-1)w_2^{k-1}\sum_{i=0}^{w_1-1}q^{w_2i}B_{k,q^{w_1w_2}}(y_1+\frac{i}{w_1})\\
&=w_2^{n-1}\sum_{k=0}^{n}\binom{n}{k}S_{n-k,q^{w_2}}(w_1-1)w_1^{k-1}\sum_{i=0}^{w_2-1}q^{w_1i}B_{k,q^{w_1w_2}}(y_1+\frac{i}{w_2}).
\end{aligned}
\tag{4.20}
$$

Letting further $w_2 = 1$ in (4.20), we get the following corollary. This is the multiplication formula for q-Bernoulli polynomials (cf.[5,(2.26)]) together with the relatively new identity mentioned in (4.18).

Corollary 4.13. *Let w_1 be any positive integer.*

$$
\begin{aligned}
B_{n,q}(w_1y_1)&=w_1^{n-1}\sum_{i=0}^{w_1-1}q^{i}B_{n,q^{w_1}}(y_1+\frac{i}{w_1})\\
&=\sum_{k=0}^{n}\binom{n}{k}B_{k,q^{w_1}}(y_1)S_{n-k,q}(w_1-1)w_1^{k-1}.
\end{aligned}
$$

Theorem 4.14. *Let w_1, w_2, w_3 be any positive integers. Then we have the following three symmetries in w_1, w_2, w_3:*

$$\begin{aligned}
&(w_1w_2)^{n-1}\sum_{i=0}^{w_1-1}\sum_{j=0}^{w_2-1} q^{w_2w_3i+w_1w_3j}B_{n,q^{w_1w_2}}(w_3y_1+\frac{w_3}{w_1}i+\frac{w_3}{w_2}j)\\
=&(w_2w_3)^{n-1}\sum_{i=0}^{w_2-1}\sum_{j=0}^{w_3-1} q^{w_1w_3i+w_1w_2j}B_{n,q^{w_2w_3}}(w_1y_1+\frac{w_1}{w_2}i+\frac{w_1}{w_3}j)\\
=&(w_3w_1)^{n-1}\sum_{i=0}^{w_3-1}\sum_{j=0}^{w_1-1} q^{w_1w_2i+w_2w_3j}B_{n,q^{w_1w_3}}(w_2y_1+\frac{w_2}{w_3}i+\frac{w_2}{w_1}j).
\end{aligned}\tag{4.21}$$

Letting $w_3 = 1$ in (4.21), we have the following corollary.

Corollary 4.15. *Let w_1, w_2 be any positive integers.*

$$\begin{aligned}
&w_1^{n-1}\sum_{j=0}^{w_1-1} q^{w_2j}B_{n,q^{w_1}}(w_2y_1+\frac{w_2}{w_1}j)\\
=&w_2^{n-1}\sum_{i=0}^{w_2-1} q^{w_1i}B_{n,q^{w_2}}(w_1y_1+\frac{w_1}{w_2}i)\\
=&(w_1w_2)^{n-1}\sum_{i=0}^{w_1-1}\sum_{j=0}^{w_2-1} q^{w_2i+w_1j}B_{n,q^{w_1w_2}}(y_1+\frac{i}{w_1}+\frac{j}{w_2}).
\end{aligned}\tag{4.22}$$

Theorem 4.16. *Let w_1, w_2, w_3 be any positive integers. Then we have the following two symmetries in w_1, w_2, w_3:*

$$\begin{aligned}
&\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,q^{w_3}}(w_1y)B_{l,q^{w_1}}(w_2y)B_{m,q^{w_2}}(w_3y)w_3^kw_1^lw_2^m\\
=&\sum_{k+l+m=n}\binom{n}{k,l,m}B_{k,q^{w_2}}(w_1y)B_{l,q^{w_1}}(w_3y)B_{m,q^{w_3}}(w_2y)w_2^kw_1^lw_3^m.
\end{aligned}\tag{4.23}$$

Theorem 4.17. *Let w_1, w_2, w_3 be any positive integers. Then we have the following two symmetries in w_1, w_2, w_3:*

(4.24)

$$\sum_{k+l+m=n}\binom{n}{k,l,m}S_{k,q^{w_3}}(w_1-1)S_{l,q^{w_1}}(w_2-1)S_{m,q^{w_2}}(w_3-1)w_3^{k-1}w_1^{l-1}w_2^{m-1}$$

(4.25)

$$=\sum_{k+l+m=n}\binom{n}{k,l,m}S_{k,q^{w_2}}(w_1-1)S_{l,q^{w_1}}(w_3-1)S_{m,q^{w_3}}(w_2-1)w_2^{k-1}w_1^{l-1}w_3^{m-1}.$$

Putting $w_3 = 1$ in (4.24) and (4.25) and mulitplying the resulting identity by w_1w_2, we get the following corollary.

Corollary 4.18. *Let w_1, w_2 be any positive integers.*

$$\sum_{k=0}^{n}\binom{n}{k}S_{k,q^{w_1}}(w_2-1)S_{n-k,q}(w_1-1)w_1^k$$
$$=\sum_{k=0}^{n}\binom{n}{k}S_{k,q^{w_2}}(w_1-1)S_{n-k,q}(w_2-1)w_2^k.$$

REFERENCES

1. E. Deeba and D. Rodriguez, *Stirling's and Bernoulli numbers* , Amer. Math. Monthly . **98** (1991), 423-426.
2. F. T. Howard, *Applications of a recurrence for the Bernoulli numbers,* , J. Number Theory **52**(1995), 157-172.
3. D. S. Kim and K. H. Park, *Identities of symmetry for Bernoulli polynomials arising from quotients of Volkenborn integrals invariant under S_3*, submitted.
4. T. Kim, *Symmetry p-adic invariant integral on $\mathbb{Z}_p$ for Bernoulli and Euler polynomials,* J. Difference Equ. Appl. **14** (2008), 1267-1277.
5. T. Kim, *On the symmetries of the q-Bernoulli polynomials* , Abstr. Appl. Anal. **2008**(2008), 7 pages(Article ID 914367).
6. W. H. Schikhof, *Ultrametric calculus: An introduction to p-adic analysis,* , Cambridge University Press, 2006.
7. Y. Simsek, *Twisted (h,q)-Bernoulli numbers and polynomials related to twisted (h,q)-zeta function and L-function*, J. Math. Anal. Appl. **324** (2006), 790-804.
8. H. Tuenter, *A symmetry of power sum polynomials and Bernoulli numbers,* Amer. Math. Monthly **108** (2001), 258-261.
9. S. Yang, *An identity of symmetry for the Bernoulli polynomials,* Discrete Math. **308** (2008), 550-554.

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 121-742, KOREA
E-mail address: `dskim@sogong.ac.kr`